



Peaks In Emission Lines In The Spectra Of Quasars

¹Y.P. Varshni, ¹John Talbot, ²Zhengo Ma

¹Physics Dept. Univ. of Ottawa, ²Nat. Astron. Obs., Chinese Academy of Science



Wavelength peaks compared with Wolf-Rayet stars		
No.	Peak (Å)	Reported wavelengths in Wolf-Rayet (Å)
1	3356	3358.6 Underhill (1959). N III λ 3355, O III λ 3355.9, C III λ 3358
2	3489	3493 Wright (1918). O IV λ 3490.8
3	3526	not reported in WR
4	3549	not reported in WR.
5	3610	3611 Wright (1918), 3609.5 Beals (1930), 3609+ Edlén (1956), 3608.5 Underhill (1959), C III λ 3609.6, He I λ 3613.6.
6	3648	3645.4 Underhill (1959). C III, O IV λ 3642
7	3683	3687 Edlén (1956), 3685.10 - novae-like stars (Meinel et al., 1975). C IV 3722 Plaskett (1924), 3723 Beals (1930), 3722 Edlén (1956), 3717.1 Underhill (1959). O III λ 3721
8	3719	3769 Plaskett (1924), 3769 Edlén (1956), 3770.6 Underhill (1962). O III, N III λ 3773
9	3770	3781 3784.8 Underhill (1959). He II λ 3781.68, O III, N III λ 3779
10	3781	3829.9 Underhill (1959). He II λ 3833.80, N IV, O VI
11	3831	Not seen in WR. O IV λ 3841.07(?), C III λ 3844.51 (?)
12	3842	3856.6 Underhill (1959). He II λ 3858.07
13	3855	3889 Wright (1918), 3889 Plaskett (1924), 3888.9 Beals (1930), 3888, 3888.7 Swings (1942), 3887.8, 3890.9 Underhill (1959), 3889.4 Underhill (1962). He I λ 3888.64, C III λ 3889.18, 3885.99
14	3890	3903.0 - novae-like stars (Meinel et al., 1975)
15	3903	3953.7 Beals (1930), 3954.4 Edlén (1956), 3954.5 Underhill (1962). O II, (C II)
16	3952	4008.2 Underhill (1959), 4008.5 Underhill (1962). N III λ 4007.88, 4013.00
17	4012	O II λ 4132.8, Possible in WR stars Edlén (1956)
18	4135	4275.5 novae (Meinel et al., 1975), 4276.6 novae-like stars (Meinel et al., 1975). O II λ 4275.5 Possible in WR (Edlén, 1956)
19	4276	4519.5 Plaskett (1924) 4521.3 Underhill (1959). N III λ 4523.56, 4527.9, O III λ 4524.2 4527.3, C III
20	4524	4650.8 Swings (1942). C III λ 4647.40, 4650.16, 4651.35; O II λ 4649.15
21	4647	4697.0 novae-like stars (Meinel et al., 1975). O II λ 4596.2 Possible in WR stars (Edlén, 1956)
22	4693	4772.1 novae-like stars (Meinel et al., 1975). O IV λ 4772.6 Possible in WR stars (Edlén, 1956)
23	4771	4799 Wright (1918), 4800 Plaskett (1924), 4798.3 Edlén (1956), 4797.4 Underhill (1959), 4798.1 Underhill (1962), 4804.6 Underhill (1962). O IV λ 4801
24	4801	4814.6 Underhill (1962), 4814.4 novae-like stars (Meinel et al., 1975). O IV λ 4813 4824, Si III
25	4817	4909.2 Underhill (1959). N III ?
26	4910	4923 Wright (1918), 4924 Plaskett (1924), 4924 Edlén (1956), 4927.4 Underhill (1959). He I λ 4921.9
27	4925	4958 Plaskett (1924), 4959.0 novae (Meinel et al., 1975), 4959.0 old novae (Meinel et al., 1975)
28	4956	5021 Campbell (1894), 5017 Wright (1918), 5018.3 Plaskett (1924), 5018 Beals (1930), 5015.7 Swings (1942), 5018 Edlén (1956), 5019.8 Underhill (1959). He I λ 5015.67, C IV λ 5015.9, 5017.7
29	5018	not seen in WR
30	5035	5049.9 Underhill (1959). He I λ 5047.7, C II
31	5049	5092.9 Swings (1942)
32	5096	5111.5 novae-like stars (Meinel et al., 1975)
33	5111	5171.1 Underhill (1959). N II λ 5172
34	5173	5266.3 Underhill (1959). C III, O III λ 5268.1
35	5266	5343.3 Swings (1942). C II λ 5336.7
36	5345	5470 Wright (1918), 5470 Beals (1930), 5470 Edlén (1956), 5469.6 Underhill (1959). C IV, O V
37	5466	

Introduction

We are startled to discover 37 very strong peaks in the distribution of emission lines (in the observed frame) in the spectra of quasars. We are further surprised to find 27 of these 37 lines in the spectra of Wolf-Rayet stars. An additional 5 lines are seen in novae like stars.




Method

We convert to observed frame 14277 rest frame emission lines listed in the 1993 Hewitt and Burbidge quasar catalog. We plot a histogram of the emission line frequency against the wavelength (middle plot below). We locate the strongest peaks by thresholding with a statistical significance of 4 standard deviations from what would be expected by chance (bottom plot)

Discussion

In the redshift hypothesis there is no reason why the emission lines in the observed frame should show these peaks. Thus the redshift hypothesis is unable to account for these peaks.

Theoretical and experimental investigations in physics in the 1960's and 1970's showed that when a high temperature plasma rapidly expands (for example, in vacuum) the resulting cooling leads to a population inversion in the lower levels of the atom, and this can lead to laser action



Conclusion: Plasma-Laser Star Model

This led Varshni (1975, ApSS 37, L1; 1977, ApSS 46, 443; 1979 Phys. Canada 35, 11) to propose that a quasar is a star in which the surface plasma is undergoing rapid radial expansion giving rise to population inversion and laser action in some of the atomic species. The assumption of high speed matter ejections from quasars is supported from the fact that the widths of emission spectral lines observed in quasars are typically of the order of 2000-4000 km/sec. Ejected matter either forms a nebulosity around the quasar or dissipates into space. Laser action is enhanced if hot plasma contacts this colder gas. No redshifts are needed. It is known that some atomic transitions are more susceptible to laser action than others. The peaks correspond to such transitions and such lines occur more frequently in quasar spectra.

